

# Medication Schedule: Outpatient Fludarabine/Melphalan with Post-Transplant Cytoxan® (PTCy) Chemotherapy

Bring this sheet with you to every visit. Starting 3 days before your transplant, bring this sheet and your medications with you to every visit.

5 days before your transplant	Date:
-------------------------------	-------

### Medications you will get in infusion:

- Chemotherapy medication: Fludarabine (Fludara®) 30 mg/m<sup>2</sup>
- Anti-nausea and anti-vomiting medications: Palonosetron (Aloxi®), prochlorperazine (Compazine®), and some patients may also get dexamethasone (Decadron®)

# Medications you may take at home (write down any time you take the medication):

Medication	Dose	Time
Prochlorperazine (Compazine®)		
Take as needed every 6 hours		
for nausea/vomiting		

4 days before your transplant	Date:
-------------------------------	-------

#### Medications you will get in infusion:

- Chemotherapy medication: Fludarabine (Fludara®) 30 mg/m<sup>2</sup>
- Anti-nausea and anti-vomiting medications: Prochlorperazine (Compazine®), and some patients may also get dexamethasone (Decadron®)

**Medications you may take at home** (write down any time you take the medication):

Medication	Dose	Time
Prochlorperazine (Compazine®)		
Take as needed every 6 hours		
for nausea/vomiting		

3 days before your transplant	Date:
-------------------------------	-------

#### Medications you will get in infusion:

- Chemotherapy medication: Fludarabine (Fludara®) 30 mg/m<sup>2</sup>
- Anti-nausea and anti-vomiting medications: Prochlorperazine (Compazine®), and some patients may also get dexamethasone (Decadron®)

Med	ications	you will	I take a	t home	(write	down	any	time	you t	ake t	the	med	icat	ion)	):
-----	----------	----------	----------	--------	--------	------	-----	------	-------	-------	-----	-----	------	------	----

Medication	Dose	Time
Prochlorperazine (Compazine®)		
• Take as needed every 6		
hours for nausea/vomiting		

2 days before your transplant	Date:
-------------------------------	-------

#### Medications you will get in infusion:

- Chemotherapy medication: Fludarabine (Fludara®) 30 mg/m²
- Anti-nausea and anti-vomiting medications: Palonosetron (Aloxi®), prochlorperazine (Compazine®), and some patients may also get dexamethasone (Decadron®)

# **Medications you will take at home** (write down any time you take the medication):

Medication	Dose	Time
Prochlorperazine (Compazine®)		
• Take as needed every 6 hours		
for nausea/vomiting		

1 day before your transplant	Date:
------------------------------	-------

## Medications you will get in infusion:

• Chemotherapy medication: Melphalan140 mg/m²

Blood and Marrow Transplant (BMT) Program

Medication Schedule: Outpatient Fludarabine/Melphalan with Post-Transplant Cytoxan® (PTCy) Chemotherapy

 Anti-nausea and anti-vomiting medications: Prochlorperazine (Compazine®), and some patients may also get dexamethasone (Decadron®)

**Medications you will take at home** (write down any time you take the medication):

Medication	Dose	Time
Prochlorperazine (Compazine®)		
Take as needed every 6 hours		
for nausea/vomiting		

Disclaimer: This document contains information and/or instructional materials developed by University of Michigan (U-M) Health for the typical patient with your condition. It may include links to online content that was not created by U-M Health and for which U-M Health does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Authors: Tate Feeney, Denise Mouro Edited by: Brittany Batell, MPH MSW

Patient Education by <u>U-M Health</u> is licensed under a <u>Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License. Last Revised 09/2023</u>