

Exercise Instructions: Carotid Artery Disease

Always follow the advice of your provider (doctor/nurse practitioner/physician assistant). The instructions below are general and should only be done with their approval.

General Guidelines

Warm-up	5 minutes
Aerobic	30-60 minutes continuous exercise (or as tolerated)
Weight Training	10-20 minutes
Cool Down	5 minutes
Stretching	Up to 5 minutes of light stretching

Warm Up

Easy/light movement that prepares muscles for aerobic and/or weight training exercise.

Aerobic

- Done most days of the week. Goal is exercise time not intensity.
- Uses large muscle groups working together.
- This should cause an increase in heart rate (HR) and breathing rate, but you should still be able to carry on a conversation.
- Examples: walking, swimming, biking, and using a Nustep®.

If unable to perform 20 minutes of continuous exercise, try interval training.

- Continuous exercise - exercise at the same pace for the entire time

- Interval training - vary exercise with easier to harder segments, for example:
 - Walk at a comfortable pace for 3 minutes,
 - Then try increasing speed or elevation for up to 1 minute,
 - Then return back to original pace for another 3 minutes,
 - Repeat as you are able

Weight Training

- Aim for high repetitions (reps) with light weights 2-3 days per week.
- Start with 1 set of 15 reps per muscle group (legs, back, chest, shoulders, and arms).
- Increase up to 3 sets of 15 reps when able to tolerate.
- Strength training should only be performed after some form of aerobic warm-up.
- Be sure to breathe out on the exertional part of the exercise (when you are contracting the specific muscle you are training).

Note: It is not safe to do isometric training (pushing or pulling against a fixed object like a wall or bar) at this time.

Cool Down

Similar to the warm up; light movement to help bring your heart rate (HR) back to its resting level.

Stretching

- Decreases muscle soreness and increases flexibility.
- Done after every workout session.
- Hold stretch up to 30 seconds (as tolerated) without moving, and be sure to keep breathing.

PRECAUTIONS

- Your physician should clear you before you start an exercise program.
- Exercise may result in fluid loss (dehydration) so drink fluids to prevent low blood pressure after exercise. Patients on fluid restriction should ask for more specific advice from their physician.

WHEN TO CALL & WHO TO CALL

WARNING SIGNS OF STROKE are a main concern when exercising. Stop all exercise and call your physician if you experience:

- Blurred vision in one or both eyes
- Weakness/numbness in your legs, arms or face on one side of your body
- Slurred speech
- Inability to understand conversation around you
- Loss of coordination
- Dizziness/confusion
- Difficulty swallowing
- Sudden onset of headache with unknown cause
- Chest pain
- Shortness of breath

BORG RATING OF PERCEIVED EXERTION (RPE)

Use this scale to find the intensity level of your exercise. A rating of 11-13 is ideal.

6	
7	Very, Very Light
8	
9	Very Light
10	
11	Fairly Light
12	
13	Somewhat Hard
14	
15	Hard
16	
17	Very Hard
18	
19	Very, Very Hard
20	

For questions about exercise, call the exercise consultants at:

Cardiovascular Center 734-936-5627

Cardiovascular Medicine at Domino's Farms 734-998-5666

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Developed by Preventive Cardiology

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](#).

Last Revised 4/2018