

Treatment Options For Early Pregnancy Loss Or Miscarriage

What are my treatment options?

You have several options to treat your early pregnancy loss or miscarriage. All of these options are safe and complications are rare. Some options are more likely than others to require more than one treatment. The treatment options differ in the length of time it takes to completely pass the pregnancy. Sometimes your provider will recommend or discourage treatments based on your medical or pregnancy history. However, most of the time, you may choose which option you most prefer.

Can I wait for the pregnancy to pass on it's own?

Often, you may wait (expectant management) for the pregnancy to pass on its own. Eventually, your uterus will begin to contract and will pass the pregnancy tissue. This can take several weeks or longer and is successful roughly 50% of the time. Heavy bleeding and painful cramping is normal with this process. Your provider will give you guidelines to watch your bleeding during this process. Risks include very heavy bleeding or incomplete passage of the pregnancy that can require surgery and infection.

Is there medicine I can take to pass the pregnancy?

You may use a medicine called misoprostol that will cause your uterus to contract and pass the pregnancy. This is successful about 85% of the time within 1 week of taking it. Heavy bleeding and painful cramping is normal with this process. If the pregnancy does not pass, additional doses of misoprostol or a surgical procedure may be needed. Risks include very heavy bleeding or incomplete passage of the pregnancy that can require surgery and possible infection. Neither of these risks is common.

Is there a surgical procedure that can pass the pregnancy?

You may have a surgical procedure to pass the pregnancy from your uterus called uterine aspiration. Your cervix (entrance to the uterus) is slowly stretched open to allow a small plastic tube to enter. Suction is applied to the tube as it is moved inside the uterus for 2-3 minutes. You will feel cramping during and after this procedure. The doctor will check to make sure that the pregnancy tissue appears completely removed. This procedure can be done in the office or in the operating room. Your provider may recommend one location over the other because of your medical history, but most of the time you may choose.

Uterine aspiration in the office

You will be awake for the procedure. You can have either pills, or medications through an IV line to assist you with pain or nervous feelings. A support person can join you in the office for the procedure. This is typically a short visit (1 hour or less).

Uterine aspiration in the operating room

You can choose to be completely asleep for the procedure. A support person will not be able to join you in the operating room for the procedure. You will spend several hours at the hospital for this procedure.

Risks include infection (up to 3 out of 100 women), incomplete aspiration (up to 3 out of 100 women), serious bleeding (1 out of 1000 women), injury to uterus or cervix (1 out of 500 women) or injury to a nearby organ (very rare).

Disclaimer: This document contains information and/or instructional materials developed by the University of Michigan Health System (UMHS) for the typical patient with your condition. It may include links to online content that was not created by UMHS and for which UMHS does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by [University of Michigan Health System](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](#) Last Revised 6/2015