



## What is an Anorectal Manometry Test?

Anorectal Manometry measures the tone (strength) in the anal sphincter and rectal muscles. This test can diagnose problems with defecation (moving your bowels). It can measure:

- resting anal sphincter tone
- releasing anal sphincter tone
- squeezing anal sphincter tone
- muscle coordination
- expulsion of materials
- sensation of the rectum

**Anal sphincter:**  
muscles at the end of the rectum that surround the anus.

**Rectal muscles:**  
muscles in the rectum-lower part of your large intestine where your body stores stool

## What can I expect from an Anorectal Manometry?

1. You will be asked to undress from the waist down and to put on a gown with the opening in the back.
2. A trained technologist will explain everything that will happen. The technologist will ask you to lie on an examination table, turn onto your left side, and bend your knees.
3. A small catheter with a small balloon at the end is inserted into the rectum.
4. The balloon is inflated and measurements are taken while you squeeze and relax your muscles.
5. You will be asked to expel the balloon into the toilet.

This test is performed in the GI Physiology Laboratory in the Medical Procedures Unit. The appointment may take up to 1 hour.

## **How do I prepare for an Anorectal Manometry?**

One and a half hours (90 minutes) before leaving home, you will need to perform two 500 cc (about 2 cups) warm tap water enemas. You may purchase the enema bag at a pharmacy or medical supply store. The enema usually takes 15 minutes to work but may take up to an hour.

1. Fill the enema bag with 500 cc (about 2 cups) warm tap water. Do not use hot water.
2. Hang the enema bag on a nail or hook or have someone hold it about 12 to 18 inches above your rectum.
3. Lie on your left side with your knees bent toward your chest.
4. Remove the cap from the enema tip.
5. Gently insert the enema tip about 3 to 4 inches into your rectum.
6. Open the clamp and allow the water to flow slowly into your rectum. The flow rate can be adjusted by moving the clamp.
7. Hold the water in your rectum as long as possible - at least 15 minutes - then expel it into the toilet.
8. Perform the second enema after you have expelled the water from the first one.

**If you are unable to keep your appointment, please call the Medical Procedures Unit at (734) 936-9250 or toll-free at (877) 758-2626.**

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#). Last Revised 01/2019