

Laparoscopic Abdominal Surgery

Are there any diet restrictions?

- Return to your same diet as before surgery
- Taking smaller portions, more frequently during the day may help in transitioning to your regular diet (6-8 small meals per day)
- Avoid constipation; drink plenty of water and caffeine-free fluids. Take stool softener while you are taking prescribed pain medication

When can I go back to normal activities?

- No lifting anything greater than 5 - 10 pounds for 6 weeks. This includes strenuous activities such as vacuuming, lifting children, groceries, laundry, moving furniture, mowing lawns, sports activities.
- No driving until your first post-op visit. You may ride as a passenger.
- You may climb stairs in moderation.
- Walk as much as tolerated. This is the only activity/exercise allowed for 6 weeks.
- Resume sexual activity as discussed at your first post-op visit.

How will I care for my incision?

- You may shower your incisions once home. Gently wash incisions and pat dry. Leave open to the air and dry; no use of powders or creams. No tub baths, hot tubs, or swimming until the incisions are completely healed
- Avoid tight clothing around your incision sites or fabrics which may irritate the skin. Wear loose comfortable clothing
- If steri strips are present, these will fall off in about 1-2 weeks. **Do not** pull them off earlier.

How I manage my pain at home?

- For best pain control take your pain medication at regular intervals, about every 4-6 hrs. This will prevent pain build-up that may occur. Use only as much prescription pain medications as you need.
- Taking over the counter ibuprofen (Motrin®, Advil®) alternated with prescription pain medication may improve pain control
- Do not drink alcohol or drive while taking narcotic pain medication

When should I call my doctor?

Call if you have any of the signs and symptoms that may indicate an infection:

- Temperature above 101°F
- Significant increase in wound pain or discomfort
- Excessive redness, swelling, or drainage from the incision sites. It is normal to have scrotal swelling and bruising
- The incisions begin to open

Call if you have any changes in your overall health such as having:

- Nausea
- Vomiting
- Chills
- profuse (excessive) sweating
- diarrhea
- constipation
- inability to urinate or completely empty bladder

What is the number to call?

- On weekdays contact the General Surgery Call Center at 734- (734)-936-5738 (Monday - Friday, 8am -5 PM)
- On weekends, holidays, or after business hours call 734-936-6267 (hospital paging) and ask for the surgery resident on-call.

What is my follow-up care?

- You will return to the clinic for your follow-up visit about two weeks after discharge. If you have not already received an appointment, we will arrange it and notify you via the patient portal or by mail.

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](#). Last Revised 2/2017