

Maxillary Prosthesis Care Instructions

What is a maxillectomy?

A maxillectomy/ palatectomy is a surgical procedure removing all or part of the hard or soft palate. The resulting defect makes swallowing, speaking, chewing, and the use of regular denture appliances almost impossible.

What is a maxillary prosthesis?

A maxillectomy/ palatectomy prosthesis or "obturator" restores the surgical defect and aids in the function of speaking, chewing, or swallowing. It fills the void left by the surgery and artificially replaces loss of tissues and teeth.

Prosthetic treatment involves:

- Patient counseling and instruction
- Physical therapy (oral - physical)
- Dietary counseling
- Hygiene maintenance and instruction
- Fabrication of prosthetic appliances

Fabrication of obturators is usually accomplished in three phases:

- Immediate -This is the surgical prosthesis it will be placed at the time of surgery and is secured in place by either wires or small screws. It will remain in place until your 1st post op visit with the prosthodontist.
- Transitional - this phase is started 10-14 days after surgery when surgical dressings or immediate prosthesis are removed. The transitional prosthesis is placed and modified until healing is complete. This phase may extend from 2-24 months.

- Definitive - this phase begins when healing is complete and involves fabrication of prosthesis intended for long term use. Definitive treatment may involve fixed prosthesis (crowns) and / or removable prosthesis.

Any phase of treatment may be altered dependant on the nature of the disease and its staging, radiation, chemotherapy, surgical complications, and the morbidity of the disease.

Obturator Care

- Clean the obturator after meals and at bedtime. Use soap and water or denture paste on a brush to remove plaque. Stains can be removed with commercial soaking agents.
- Clean the obturator over 4-5 inches of water in a basin or sink to avoid damage to the obturator should it drop.
- Remove the prosthesis at night and place it in a container of water. (Please note, that the surgical obturator is not to be removed).

Who should I call if I have questions?

- (734) 936-5950, Monday - Friday, 8 a.m. - 4:30 p.m.
- After hours and on weekends, call Hospital Paging at (734) 936-6267 and ask for the Oral & Maxillofacial Surgeon on call.

Disclaimer: This document contains information and/or instructional materials developed by the University of Michigan Health System (UMHS) for the typical patient with your condition. It may include links to online content that was not created by UMHS and for which UMHS does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Author: Erin Larowe
Reviewers: Brent Ward, DDS, MD

Patient Education by [University of Michigan Health System](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](#). Last Revised 3/2016