

艾多沙班 (Savaysa®)

Edoxaban (Savaysa®)

什么是艾多沙班 (Savaysa®)?

What is edoxaban (Savaysa®)?



- 艾多艾多沙班是个血液稀释药物。 这药物的商品名也称为 Savaysa®.
- Edoxaban is a blood thinner medication. This medication is also called by its brand name, Savaysa®.
- 艾多沙班 (Savaysa®) 用于降低患有异常心律 (称为心房纤颤) 的人发生血栓和中风的风险。
 - Edoxaban (Savaysa®) is used to reduce the risk of blood clots and stroke in people with an abnormal heart rhythm (called atrial fibrillation).
 - 血栓能堵塞血管，切断该区域的血液供应。
 - Blood clots can block a blood vessel, cutting off blood supply to the area.
 - 在极少数的情况下，血栓会破裂成片并进入血液循环，这些碎片可能会卡在心脏 (导致心脏病发作) 、肺部 (导致肺栓塞) 或脑部 (导致中风) 。
 - Rarely, clots can break into pieces and travel in the bloodstream.

Those pieces may get stuck in the heart (causing a heart attack), the

lungs (causing a pulmonary embolus), or in the brain (causing a stroke).

- 如果您之前在用华法林 (warfarin) 且现在开始用艾多沙班 (Savaysa®)，**请勿继续使用华法林 (Coumadin®)**。艾多沙班 (Savaysa®) 是替代华法林/Coumadin®的。
- If you were taking warfarin (Coumadin®) and you are starting to take edoxaban (Savaysa®), do not continue taking warfarin. Edoxaban (Savaysa®) replaces warfarin.

我应该如何服用艾多沙班 (Savaysa®)?

How should I take edoxaban (Savaysa®)?

- 请准确按照医生规定的方式服用艾多沙班 (Savaysa®)。
- Take edoxaban (Savaysa®) exactly as prescribed by your doctor.
- 艾多沙班 (Savaysa®) 可与食物一起服用，亦可单独服用。
- You can take edoxaban (Savaysa®) with or without food.
- 如果您忘记服药，在当天想起此事请尽快补服。请勿为了弥补忘记服的药而服用双倍剂量。
- If you miss a dose, take the dose as soon as you remember on the same day. Do not take a double dose to make up for missing a dose.

- 请勿在未告知您医生的情况下突然停止服用艾多沙班 (Savaysa®)。否则您将可能有中风或血栓的风险。
- Do not stop taking edoxaban (Savaysa®) suddenly without telling your doctor. This can put you at risk of having a stroke or a blood clot.
- 如果您过量服用艾多沙班 (Savaysa®)，请致电您医生或抗凝诊所。如果您出现了无法止住的出血，请前往最近的急诊室。
- If you take too much edoxaban (Savaysa®), call your doctor or the Anticoagulation Clinic. If you are experiencing any bleeding which you cannot get to stop, go to the nearest emergency room (ER).

我应该如何存放艾多沙班 (Savaysa®)?

How should I store apixaban (Eliquis®)?

请将它存放于常温干燥处。 避免存放于卫生间。

Store this medication at room temperature in a dry area. Avoid storing it in the bathroom.

我该知道服用艾多沙班 (Savaysa®) 有哪些风险?

What should I know about the risks of taking edoxaban (Savaysa®)?

- 艾多沙班 (Savaysa®) 能增加您出血的风险，因为它可减弱您身体的凝血功能。在服用艾多沙班的期间，您可能更容易瘀伤并可能需要更长时间才能止血。
- Edoxaban (Savaysa®) can increase your risk of bleeding because it reduces your blood's ability to clot. While you take edoxaban (Savaysa®), you may bruise more easily and it may take longer for bleeding to stop.
- 如果您服用艾多沙班 (Savaysa®)，且符合以下情况，则您出血的风险可能更高：
- You may have a higher risk of bleeding if you take edoxaban (Savaysa®) and:
 - 年满 75 岁 以上
 - You are over 75 years old.
 - 患有肾脏疾病
 - You have kidney problems.
 - 最近出现或复发(消失了然后又出现)胃或肠道出血
 - You have stomach or intestine bleeding that is recent or recurs (goes away and then comes back).
 - 或患有胃溃疡
 - You have a stomach ulcer.
 - 您服用其他药物可能导致出血几率升高，包括：
 - You take other medications that increase your risk of bleeding, including:
 - 阿司匹林或含阿司匹林的药物。

- Aspirin or products that contain aspirin
- 长期服用非甾体抗炎药 (Non-steroidal Anti-inflammatory Drugs, NSAIDs), 包括布洛芬及萘普生。
- Long-term use of non-steroidal anti-inflammatory drugs (NSAIDs), including ibuprofen and naproxen
- 氯吡格雷 (Plavix®)、普拉格雷 (Effient®) 或替格瑞洛 (Brilinta®)
- Clopidogrel (Plavix®), prasugrel (Effient®), or ticagrelor (Brilinta®)
- 所有含肝素的药物
- Any medication that contains heparin

艾多沙班 (Savaysa®)有哪些潜在的副作用?

What are the possible side effects of edoxaban (Savaysa®)?

- 艾多沙班 (Savaysa®) 能增加您出血的风险，因为它可减弱您身体的凝血功能。在服用艾多沙班的期间，您可能更容易瘀血并需要更长时间才能止血。
- Edoxaban (Savaysa®) can increase your risk of bleeding because it reduces your blood's ability to clot. While you take edoxaban (Savaysa®), you may bruise more easily and it may take longer for bleeding to stop.
- 部分人群服用艾多沙班 (Savaysa®)后可能出现过敏反应，包括荨麻疹、皮疹和瘙痒。如果您出现以下任一严重过敏反应，请立即寻求医疗救助：

- For some people, edoxaban (Savaysa®) can cause symptoms of an allergic reaction, including hives, rash, and itching. Get medical help immediately if you get any of these symptoms of a serious allergic reaction:
 - 胸痛或胸闷
 - Chest pain or chest tightness
 - 面部或舌头肿胀
 - Swelling of your face or tongue
 - 呼吸困难或气喘
 - Trouble breathing, or wheezing
 - 感到头晕或即将晕厥
 - Feeling dizzy or faint

我应该何时致电我的医生或抗凝诊所?

When should I call my doctor or the Anticoagulation Clinic?

如果有下列状况, 打电话给您的医生或打给我们

Call your doctor, or call us, if:

- 在您开始或停止使用任何药物的时候 (包括非处方药物或草药补品) 。因为它可能影响艾多沙班 (Savaysa®) 的服用效果。
- You start or stop any medications, including over-the-counter medications or herbal supplements. These may affect the way edoxaban (Savaysa®) works.

- 您将进行外科手术或其他医疗程序时，因为您可能会需要了解如何服用艾多沙班 (Savaysa®)的特别指示。
- You are having a surgery or a procedure, as you may have special instructions on how to take your edoxaban (Savaysa®).
- 在您需要补购药物的时候。在您快要服用完艾多沙班 (Savaysa®) 之前告知您的医生，以避免药量不足）。
- You need a medication refill. Try to tell your doctor before you are out of edoxaban (Savaysa®) so you do not run out.
- 在您意外出血或长时间出血的时候，例如：
- You have any unexpected bleeding or bleeding that lasts a long time, such as:
 - 牙龈异常出血
 - Unusual bleeding from the gums
 - 频繁流鼻血
 - Nose bleeds that happen often
 - 月经(经期)或阴道出血量异常多
 - Menstrual (period) or vaginal bleeding that is heavier than normal
 - 尿中有血(尿液呈红色或咖啡或可乐的颜色)
 - Blood in your urine (pee that is red or the color of coffee or cola)
 - 出现未知原因的瘀血或瘀血范围增大

- Bruises that happen without a known cause, or bruises that get larger

我应该何时寻求紧急医疗救助？

When should I get emergency help?

如果您出现以下任一严重情况，请立即致电 911：

Call 911 immediately if you have any of the following serious conditions:

- 如果您出现以下任何严重出血的体征和症状：
 - 出血情况严重或您无法止血
 - Severe bleeding, or bleeding you can't control
 - 大便呈红色或黑色（大便如沥青一样）
 - Red or black stools (poop that looks like tar)
 - 咳血或咳血块
 - Coughing up blood or blood clots
 - 呕吐出血或您的呕吐物看起来像咖啡渣
 - Vomiting up blood, or if your vomit looks like coffee grounds
 - 鼻子或牙龈流血不止
 - Uncontrolled bleeding from your nose or gums
 - 割伤不能止血

- Cuts that will not stop bleeding
- 如果您出现以下任一中风体征和症状:
 - 头晕、突发性视力丧失或突发性严重头疼
 - Dizziness, sudden loss of vision, or a sudden severe headache
 - 任何一个手臂或腿麻木或无力
 - Numbness or weakness in any arm or leg
 - 说话困难或口齿不清
 - Trouble speaking, or slurred speech
 - 面部一侧下垂
 - Drooping on one side of your face
- 如果您出现以下任一血栓体征和症状:
 - You have signs or symptoms of blood clots:
 - 突发性严重呼吸短促
 - Sudden severe shortness of breath
 - 胸痛
 - Chest pain
 - 任何一个手臂或腿发生肿胀、泛红、发热或疼痛
 - Swelling, redness, heat or pain in any arm or leg

免责声明：本文件所包含的信息和/或教学材料是由密西根大学健康中心为与您病情类似的典型患者所撰写的。文中的链接可能连接到并非由密西根大学健康中心所创建的网络内容，密西根大学健康中心对此内容不承担责任。本文件不可取代您的医疗保健提供者的医疗咨询，因为您的情况可能和典型患者有所不同。

如果您对此文件、您的病情或治疗方案有任何疑问，请与您的医疗保健提供者商谈。

Disclaimer: This document contains information and/or instructional materials developed by University of Michigan (U-M) Health for the typical patient with your condition. It may include links to online content that was not created by U-M Health and for which U-M Health does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

作者: Elizabeth Renner, PharmD

审稿员: Nghi Ha, PharmD

编辑: Brittany Batell, MPH MSW

Author: Elizabeth Renner, PharmD

Reviewer: Nghi Ha, PharmD

Edited by: Brittany Batell, MPH MSW

翻译: 密西根医学部翻译服务

Translation: Michigan Medicine Interpreter Services

密西根大学健康中心病患教育是由 [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#) 授权许可。最后修订日期 05/2024

Patient Education by University of Michigan Health is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#). Last Revised 05/2024