

Port or Catheter Removal Instructions

What will happen after my procedure?

- You received local anesthesia. As the anesthesia wears off, you may feel some pain and discomfort.
- The stitches will dissolve in 10-14 days.
- The steri strips (white paper tape over the incision) will fall off in approximately 1 week.

How will I care for myself at home?

- You may resume showering in 8 hours.
 - Do not swim, use a hot tub, or take a bath for 14 days after your procedure.
- No strenuous physical activity for 2 days.
- Avoid activities that cause excessive tension or stretching across the healing wound site for 1 week.
- Resume your daily prescription medications.
- Resume your previous diet.

How do I take care of my bandage?

Keep your bandage in place for 48-72 hours. If the dressing becomes wet or soiled before then, you may change it:

1. Remove dressing
2. Wash incision with mild soap and water; pat dry
3. Apply a sterile dressing (such as a Band-Aid) over wound

After your first dressing is removed, wash daily:

1. Wash the incision site with liquid antibacterial soap
2. Rinse
3. Pat dry
4. Cover with a clean Band Aid if necessary to prevent abrasion.
5. After 1 week, no Band Aid is necessary; keep area clean and dry.

How do I care for a skin adhesive?

You may have a type a skin adhesive (glue) instead of a bandage. Do not bandage a wound treated with a skin adhesive. The skin adhesive works like a bandage.

- Do not use antibiotic ointment as it can break down the adhesive.
- You can shower while the adhesive is on your skin (same day is ok), but do not take a bath or soak or scrub the area for 14 days. Dry your skin by patting it gently with a towel.
- The adhesive will peel off on its own, usually in 5 to 10 days.
 - If it has been 10 days and you still have adhesive on you, you can use antibiotic ointment or petroleum jelly to gently rub it off.
- **Call if you have signs of infection**, such as redness, swelling, or pus.

What signs and symptoms should I expect?

Tenderness at site, minimal swelling, bruising or slight bleeding may occur.

When should I call for help?

Call Interventional Radiology if you notice any of the following:

- fever (higher than 100.5F)
- redness
- swelling
- red streaking
- pain

- pus
- excessive or prolonged bleeding

What is the contact information?

If any of the above symptoms occur:

- Monday-Friday 8 a.m.-5 p.m. call Interventional Radiology:
 - (734) 936-4500, option 2, 5
- After hours, weekends or holidays call Interventional Radiology and ask the operator for the “Interventional Radiology On-Call Fellow/Resident”:
 - (734) 936-6267
- If you cannot reach medical assistance, go to an Emergency Room. Take this form with you.

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](#). Last Revised 1/2023