

Instructions After Allergic Reaction to Contrast During Magnetic Resonance Imaging (MRI) Scan

What are my Care Instructions?

You had an allergic reaction to MRI Contrast. You might still be itchy or have hives or a rash. You might be drowsy from the medication you received.

Follow-up care is the key part of your treatment and safety. Be sure to make and go to all appointments and call your doctor if you have problems.

How can I care for myself at home?

Activity

For your safety, avoid activities that require motor coordination (such as driving) if you have received any medicines for your contrast reaction including:

- diphenhydramine (Benadryl)
- epinephrine (Auvi-Q, EpiPen)

Diet

- Eat your normal diet. If your stomach is upset, try bland, low-fat foods like plain rice, toast, and yogurt.
- Drink plenty of fluids (unless your doctor tells you not to).
- Do not drink alcoholic beverages for 24 hours.

Medications

• Take your normal medicines your doctor has prescribed.

Special Instructions

- If you need an IV contrast agent for an MRI scan in the future, please tell the Radiology Department about this reaction.
- Talk with your doctor before your next MRI scan requiring a contrast agent so they can consider preparing you with steroids and Benadryl.

Department of Radiology

When should you call for help?

Call 911 anytime you think you need emergency care. Call if you have any of the following signs and symptoms:

- Shortness of breath
- Chest pains
- Loss of consciousness (if you passed out or fainted)

Call the Radiology Nursing Station at 734-936-4515 between 8am and 5pm Monday through Friday (except holidays) if you have any of the following signs and symptoms:

- More hives or rash
- Increasing itching or troublesome itching that won't go away
- Swelling
- Dizziness
- Vomiting that does not stop
- Redness, swelling, red streaking or pain at IV site

After hours you can call the Radiology Superchief (the radiology resident on call for emergencies) at 734-936-6267, pager 1800.

Disclaimer: This document contains information and/or instructional materials developed by University of Michigan (U-M) Health for the typical patient with your condition. It may include links to online content that was not created by U-M Health and for which U-M Health does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Reviewer: James Ellis, MD

Patient Education by <u>U-M Health</u> is licensed under a <u>Creative Commons Attribution</u>-NonCommercial-ShareAlike 4.0 International Public License. Last Revised 04/2022

Department of Radiology Instructions After Allergic Reaction to Contrast During Magnetic Resonance Imaging (MRI) Scan