

To reduce your bleeding and clotting risk it is important that you **attend follow-up appointments** with your provider, and **have blood tests done** as your provider orders.

## What is Edoxaban (Savaysa®)?

- Edoxaban is also called Savaysa®
- Edoxaban(Savaysa®) is used to reduce the risk of blood clots and stroke in people with an abnormal heart rhythm known as atrial fibrillation.
  - Blood clots can block a blood vessel cutting off blood supply to the area.
  - Rarely, clots can break into pieces and travel in the blood stream, lodging in the heart (causing a heart attack), the lungs (causing a pulmonary embolus), or in the brain (causing a stroke).
- If you were previously on warfarin and you are starting Edoxaban(Savaysa®), **do not** continue taking Warfarin(Coumadin®). Edoxaban(Savaysa®) replaces Warfarin/Coumadin®

## How should I take edoxaban (Savaysa®)?

- Take Edoxaban(Savaysa®) exactly as prescribed by your doctor.
- Edoxaban(Savaysa®) may be taken with or without food.
- If you missed a dose:
  - Take it as soon as you remember on the same day. Do not double a dose to make up for a missed dose.
- Do not stop taking Edoxaban(Savaysa®) suddenly without telling your doctor. This can put you at risk of having a stroke or a blood clot.
- If you take too much Edoxaban(Savaysa®), call your doctor or anticoagulation service. If you are experiencing any bleeding which you cannot get to stop, go to the nearest emergency room.

## **How should I store edoxaban (Savaysa®)?**

- Store at room temperature in a dry area (avoid storing it in the bathroom).

## **What should I know about edoxaban (Savaysa®)?**

- Edoxaban(Savaysa®) can increase your risk of bleeding because it lessens your body's ability to clot. While you take edoxaban, you may bruise more easily and it may take longer for bleeding to stop.
- Edoxaban(Savaysa®) can cause bleeding which can be serious. You may have a higher risk of bleeding if you take Edoxaban(Savaysa®) and:
  - You are over 75 years old
  - You have kidney problems
  - You have stomach or intestine bleeding that is recent or recurs, or have a stomach ulcer.
  - You take other medications that increase your risk of bleeding, including:
    - Aspirin or products that contain aspirin
    - Long-term use of non-steroidal anti-inflammatory drugs (NSAIDs) including ibuprofen and naproxen.
    - Clopidogrel (Plavix®), prasugrel (Effient®), or ticagrelor (Brilinta®)
    - Any medicine that contains heparin

## **What are the potential side effects of edoxaban (Savaysa®)?**

- Edoxaban(Savaysa®) can increase your risk of bleeding because it lessens your body's ability to clot. While you take edoxaban, you may bruise more easily and it may take longer for bleeding to stop.
- In some people, edoxaban can cause symptoms of an allergic reaction, including hives, rash, and itching. Seek medical attention right away if you get any of the following symptoms of a serious allergic reaction:
  - Chest pain or chest tightness

- Swelling of your face or tongue
- Trouble breathing or wheezing
- Feeling dizzy or faint

### **When should I call my doctor or anticoagulation service?**

- If you start or stop any medications, including over-the-counter medications or herbal supplements. These may affect the way Edoxaban(Savaysa®) works.
- You are having a surgery or procedure as you may need to stop taking Edoxaban(Savaysa®) for a short period of time before your procedure. Do not stop taking Edoxaban(Savaysa®) without first talking with your doctor.
- You need a medication refill (notify them before you are out of Edoxaban(Savaysa®) so you do not run out).
- If you are unable to afford Edoxaban(Savaysa®)
- If you experience any unexpected bleeding or bleeding that lasts a long time, such as:
  - Unusual bleeding from the gums
  - Nose bleeds that happen often
  - Menstrual bleeding or vaginal bleeding that is heavier than normal
  - Bruises that happen without a known cause or get larger

### **When should I seek emergency medical help?**

**Call 911** immediately if you experience any of the following serious conditions:

- If you experience any of the following signs and symptoms of bleeding:
  - Bleeding that is severe or bleeding that you cannot control
  - Pink or brown urine
  - Red or black stools (that look like tar)
  - Coughing up blood or blood clots
  - Vomiting up blood or if your vomit looks like “coffee grounds”
- If you experience any of the following signs and symptoms of a stroke:
  - Dizziness, sudden loss of vision, or sudden severe headache

- Numbness or weakness in any limb
- Difficulty speaking or slurred speech
- Drooping on one side of your face
- If you experience any of the following signs and symptoms of blood clots:
  - Sudden severe shortness of breath, chest pain
  - Swelling, redness, heat, or pain in any limb

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Author: Elizabeth Renner, PharmD  
Reviewers: Nghi Ha, PharmD & Erin Mouland, PharmD

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#). Last Revised 07/2019