

How to Use Your Combivent Respimat Inhaler (ipratropium/albuterol)



What is my prescription?	Special Instructions
	Take every day to control your symptoms.

What can I expect while I take this medication?

Combivent Respimat is a short- acting anticholinergic and short-acting beta-2 agonist and will begin working within minutes. You can expect to breathe easier and it opens your small and large airways.

How to set-up your Combivent® Inhaler:

Your **Combivent Respimat** will come packaged as two parts, a cartridge and an inhaler. These pieces will need to be put together before you can use it.

1. Press the gray safety catch, while pulling off the clear base.
2. Take the cartridge out of the box and push the narrow end of the cartridge into the inhaler.

3. The cartridge can be pushed against a hard surface to make sure that it is inserted all the way.
4. About 1/8 inch of the cartridge will remain visible when the cartridge is inserted correctly.
5. Do not remove the cartridge once it has been inserted into the inhaler.
6. Put the clear base back into place.

How and when to prime your Combivent® Respimat®:

Prime inhaler if new or has not been used for more than 3 days.

1. Hold the inhaler upright with the orange cap closed.
2. Turn the clear base in the direction of the white arrows on the label until it clicks.
3. Flip the orange cap until it snaps fully open.
4. Point the inhaler toward the ground.
5. Press the dark gray button.
6. Repeat steps 1 to 5 until a spray is visible.
7. It will look like a vapor cloud.
8. After you see a visible spray; repeat steps 1 to 5 again.

If you have not used your inhaler for more than **21** days repeat steps 1 to 6.

How to use your Combivent Respimat:

1. Hold the inhaler upright with the orange cap closed.
2. Turn the clear base in the direction of the white arrows on the label until it clicks.
3. Flip the orange cap until it fully snaps open.
4. Breathe out fully (in preparation to breathe in medication).
5. Close your lips around the end of the mouthpiece without covering the holes on the side of the mouthpiece.
6. While taking in a slow, deep breath through your mouth, press the dark gray button and continue to breathe in slowly until your lungs are full.

7. Hold your breath for 5-10 seconds, and then breathe out.
8. If your physician has prescribed more than one dose (puff), wait 30 seconds and repeat above.
9. Put the cap back on.

Do not use Combivent Respimat to treat sudden symptoms of your health condition. Always carry a rescue inhaler with you.

- To view additional information about your health condition visit:
<http://www.med.umich.edu/careguides>.
- To learn more about Combivent® Respimat® visit:
www.combivent.com

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](#). Last Revised 10/2017