

Managing Acute Pain after a Dental Procedure

What pain can I expect after dental surgery?

It is normal to have some pain after dental surgery. The goal of managing your acute pain after surgery is to minimize your pain enough so that you may sleep, eat, and feel comfortable getting up and doing simple tasks. Pain is usually worst the first 3 days after your procedure and then improves. If your pain does not start to improve after 2-3 days, contact your dental provider.

What can I do to relieve pain without medications?

Rest, elevated your head, placing ice pack on your face (15 min on, 15 min off for first 24 hrs), and using warm heat can greatly reduce the amount of pain after surgery.



Distracting activities such as mindful breathing, for example music, meditation and guided imagery may be helpful to reduce pain and possibly the amount of pain medicine you will need to take to control it.

How will I use medications to relieve pain?



Research has shown that **Around-The-Clock dosing** of pain medications is very effective to control pain after surgery. This means that you will take a dose of pain-relieving medication on a set schedule rather than taking them as needed when you feel pain.

On this schedule, you will alternate pain medications acetaminophen (Tylenol®) and ibuprofen (Motrin® or Advil® and others). Some patients will also use opioid pain medication (narcotic) if their doctor prescribes it. If you have a medical condition that prohibits you from taking acetaminophen and/or ibuprofen talk with your doctor.

How may I alternate pain medication Around-The-Clock?

An example of how you may take a dose of pain medication every three hours.

- Start by taking 400-600 mg of ibuprofen (2-3 pills of 200 mg)
- 3 hours later take 650 milligram (mg) of acetaminophen (2 pills of 325 mg)
- 3 hours after that take 400-600 mg of ibuprofen
- 3 hours after taking the Ibuprofen take 650 mg of Acetaminophen

We recommend that you follow this schedule Around-The-Clock for 2-3 days after the procedure and then use the medications as needed. (Use the table on page 5 of this handout to keep track of the medications you are taking.)

What are the risks of pain medications?

- **Ibuprofen** may cause upset stomach and indigestion. Taking the medication with food or milk may help avoid these effects.
- **Acetaminophen** -taking more than 3-4 grams (3000-4000 milligrams/mg) of acetaminophen per day may cause serious injury to your liver and kidneys.

What if my doctor prescribes an opioid for me?

You may receive a prescription for opioid pain medication for break-through pain. "Break-through pain" means pain that is not controlled by Around-The-Clock acetaminophen and ibuprofen. Examples of opioids include Oxycodone, Hydrocodone, and others. **Opioids are potent medications that provide good pain relief but their side-effects may cause harm.**

If your doctor prescribes an opioid pain medication you can **replace** one of the acetaminophen or ibuprofen doses with a dose of the opioid.

For example, your schedule could look like this:

12:00 PM	Acetaminophen 650 mg (2 pills of 325 mg)
3:00 PM	Ibuprofen 600 mg (3 pills of 200 mg)
6:00 PM	Oxycodone 5mg (1 pill of 5 mg)
9:00 PM	Ibuprofen 600 mg (3 pills of 200 mg)
12:00 AM	Acetaminophen 650 mg (2 pills of 325 mg)
Continue alternating every 3 hours	

Important:

- If your pain is manageable, avoid taking opioids.
 - Never take more frequent or higher doses than prescribed
- Visit <http://michmed.org/ea93D> to learn more about taking opioids safely.

Warning: Some opioid pain medications **also contain acetaminophen** in addition to the opioid. Check the label to see if your medication contains acetaminophen and if it does make sure you **count this amount in daily intake** which is **less than 3000mg of acetaminophen in a 24-hour period** or lower (i.e. 1000mg if you have underlying medical conditions such as liver disease).

Where should I store opioid medications?

Store opioids out of sight and reach of children, teens, and pets.

- Store opioids in private areas rather than common rooms like bathrooms or kitchens.
- Lock up the pills if possible.
- Keep a count of how many pills you have left.

How should I dispose of unused opioids?

The safest way to dispose of old medications is to take them to an authorized “Take-Back” program. Visit <http://michmed.org/5LWpZ> to find a location in your area.

If you are not able to find a program in your area follow these steps as a last resort:

- Mix opioids (do not crush) with used coffee grounds or kitty litter in a plastic bag and put in your household trash.
- Scratch out personal information on the prescription label and dispose of the original container.

 **Do not flush opioids down the toilet.**

Who do I call if I have any questions or problems?

Call your dentist’s office if you have any questions, problems or if you are still having pain after 7 days.

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Modifier: Jillian Rigert, DMD, MD

Reviewer: Stephanie Munz, DDS

Plain language editing: Ruti Volk, MSI, AHIP

Modified from: Pain Management Committee. *Managing Acute Pain after Surgery*.


Access at: <http://careguides.med.umich.edu>

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 Unported License](#). Last Revised 11/28/2020

Use this table to keep track of the medications you are taking.

Share it with your doctor if your pain is not well controlled.

Use this scale to rate your pain level.



Day 1					
Time	Name of Medication	Number of pills taken	Amount of Acetaminophen (in mg)	Pain Level	Comments
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
Total Daily amount of Acetaminophen					
Do not take more than 3,000 mg per day					

Day 2					
Time	Name of Medication	Number of pills taken	Amount of Acetaminophen (in mg)	Pain Level	Comments
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
Total Daily amount of Acetaminophen					
Do not take more than 3,000 mg per day					

Day 3					
Time	Name of Medication	Number of pills taken	Amount of Acetaminophen (in mg)	Pain Level	Comments
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
Total Daily amount of Acetaminophen Do not take more than 3,000 mg per day					

Day 4					
Time	Name of Medication	Number of pills taken	Amount of Acetaminophen (in mg)	Pain Level	Comments
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
Total Daily amount of Acetaminophen Do not take more than 3,000 mg per day					

Day 5					
Time	Name of Medication	Number of pills taken	Amount of Acetaminophen (in mg)	Pain Level	Comments
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					

AM PM					
AM PM					
Total Daily amount of Acetaminophen Do not take more than 3,000 mg per day					

Day 6					
Time	Name of Medication	Number of pills taken	Amount of Acetaminophen (in mg)	Pain Level	Comments
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
Total Daily amount of Acetaminophen Do not take more than 3,000 mg per day					

Day 7					
Time	Name of Medication	Number of pills taken	Amount of Acetaminophen (in mg)	Pain Level	Comments
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
AM PM					
Total Daily amount of Acetaminophen Do not take more than 3,000 mg per day					

If your pain is not improving after 2-3 days of around the clock medications contact
Hospital Dentistry at (734)-936-5950