

Giving Midazolam from a Vial (Liquid) Intranasally

Medication: Midazolam

What do I do when I get the medication from HomeMed?

- Always check the prescription and label for final instructions.
 - check your name
 - medication name and dose
 - when the medication should be given
- If the information does not match, contact HomeMed immediately at **800-862-2731**.
- Check the vials for expired dates, color changes, leaks and floating materials. If noted, do not use the item and call HomeMed.

What supplies do I need?

Midazolam vial (5 mg per mL)

Syringe with sharp needle

Atomizer

Alcohol pads

Sharps container


How do I prepare the medication?

1. Remove the flat disk cover from the top of the liquid medication vial and **vigorously** scrub the top of the vial with an alcohol pad for fifteen seconds.
2. Remove syringe from the package and check to be sure the needle is securely attached to the syringe by twisting the needle to the right.
 - If you need to attach a sharp needle to the syringe, remove the needle from the package by holding the cap and do not touch the syringe tip as you connect it.
3. Draw the syringe plunger back to dose volume listed on the prescription label.

4. Remove the needle cap (set aside for recapping) and inject the air into the liquid vial.
5. Keeping the needle in the vial, turn the vial upside down and withdraw the dose volume listed on the prescription label. Make sure that the liquid is covering the tip of the needle.
 - If directed on medication label, draw up an additional 0.1 mL of medication to allow for dead space in the atomizer.



6. Remove the syringe from the vial and replace cap onto the needle with a one-hand scoop technique.



7. Remove the capped needle and discard in your sharps container.



8. Attach the atomizer tip to the syringe by twisting together.



How do I give the medication?

1. Using your free hand to hold the crown of the head stable, place the tip of the atomizer snugly against the nostril.
2. Push the syringe plunger and give half of the medication into the nostril.
3. Move the device over to the opposite nostril and push the syringe plunger to give the remaining medication into the opposite nostril.



What do I do if I still have questions about these instructions?

Call HomeMed. Our 24-hour number is 800-862-2731.

Please identify yourself to the hospital operator as a HomeMed patient

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Author: HomeMed Patient Education Committee

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](#). Last Revised 4/2023