

What is a Hydrogen Breath Test?

This is a test for bacterial overgrowth, or intolerance to lactose, fructose, or sucrose. Glucose is a sugar that will be broken down by bacteria (if present) in the small bowel with hydrogen or methane gas as a by-product. The breath sample will be analyzed for hydrogen or methane content to determine if you are able to properly break down the lactose, fructose, or sucrose, or if you have bacterial overgrowth.

What can I expect from a Hydrogen Breath Test?

This test requires that you drink a mixture of glucose, lactose, fructose, or sucrose in water.

- **Glucose** is a sugar that will be broken down by bacteria (if present) in the small bowel with hydrogen as a by-product.
- **Lactose** is the sugar found in milk and is normally broken down in the small bowel.
- **Sucrose** is a common sugar found in foods such as candy.

After drinking the solution, you will be asked to breathe into a plastic bag. Breath samples are obtained every 15 minutes for adults, and every 30 minutes for children. The breath sample will be analyzed for hydrogen content to determine if you are able to properly break down the lactose or sucrose, or if you have bacterial overgrowth.

The test for glucose lasts about 2 hours and the test for lactose, fructose, and sucrose lasts about 2-3 hours. The test is performed in the Gastrointestinal Physiology Laboratory of the Medical Procedure Unit.

How do I prepare for a Hydrogen Breath Test?

Please follow these instructions to ensure a successful test.

- **2 weeks before your test:**
 - You must be off antibiotics and Pepto-Bismol for 2 weeks.
- **2 days before your test:**
 - Eat a **low carbohydrate and dairy diet**. The purpose of this diet is to avoid foods that may produce gas in the intestinal tract for 2 days before your test, as this could cause inaccurate test results.

	Allowed	Avoid
Grains/starches	Rice, quinoa, potato, oatmeal, corn tortillas, popcorn, gluten-free bread/crackers/pasta. Cereals: plain cheerios, plain rice/corn Chex, corn flakes, Rice Krispies.	Wheat products including bread, pasta, crackers, cookies, cake, cereal, flour tortilla, rye barley
Vegetables	Bell peppers (any color), carrots, chives, cucumber, eggplant, green beans, kale, lettuce, olives, parsnip, spinach, summer/winter squash, zucchini, tomato. Limit to ½ cup portion: beets, butternut squash, broccoli, brussel sprouts, corn, fennel bulb, green peas, sweet potato.	Artichoke, asparagus, cauliflower, garlic, leeks, mushrooms, okra, onion/shallots, snow peas, sugar snap peas.
Fruit	Banana, blueberry, cantaloupe, cranberry, grapes, honeydew, kiwi, lemon, lime, orange, papaya, pineapple, pomegranate, raspberry, strawberry, avocado	Apple, apricot, blackberry, cherry, dried fruit (raisins), grapefruit, mango, nectarine, pear, peach, plum, prunes, watermelon.

Proteins	Meat, eggs, fish, tofu, most nuts (see “avoid” column), peanut butter, almond butter.	Beans, hummus, edamame, cashews, pistachios.
Dairy	<i>Lactose free</i> milk/yogurt/ice cream, rice milk, any aged or hard cheese (including cheddar, provolone, swiss, parmesan, feta, goat, etc)	Milk, yogurt, ice cream, cottage cheese, ricotta, cheese, soy milk, almond milk
Beverages	Water, coffee, tea, sugar-free beverages (crystal light, diet pop)	Regular soda and other beverages with high fructose corn syrup
Sweeteners	Pure maple syrup, sugar substitutes (such as Stevia), table sugar.	Agave, honey, high fructose corn syrup.

Sample Menu

Day 1

Breakfast: 2 scrambled eggs and 1 banana.

Lunch: salad with grilled chicken, tomato, cucumber, carrots, feta, balsamic vinegar + olive oil dressing.

Snack: 1 cup fruit (such as blueberries and strawberries), small handful of peanuts or almonds.

Dinner: Grilled chicken, baked potato, ½ cup steamed broccoli.

Snack: 1 cup popcorn.

Day 2

Breakfast: plain cheerios with lactose-free milk (such as Lactaid brand) or oatmeal with cinnamon and blueberries.

Lunch: sandwich with turkey, cheese, lettuce, and tomato on gluten-free bread; plain potato chips or 1 orange.

Snack: baby carrots and 1 cheese stick.

Dinner: grilled salmon with lemon, brown rice or quinoa, sautéed zucchini and yellow squash.

Snack: 1 cup grapes

- **The day of your test:**

- You should have **nothing by mouth eight 8 hours before the test.**
- **Do not eat, drink, or smoke** until your examination is completed.
- This includes any type of **gum or candy.**

Please note: breath tests can be done any time *before* a colonoscopy prep; but not for four (4) weeks *after* a colonoscopy prep, or any test that requires a bowel prep.

Can I take my medicine before a hydrogen breath test?

- Critical medicines, such as those for your heart, blood pressure, breathing, or seizures, should be taken with sips of water on the morning of the test.
- Noncritical medicines, including over-the-counter medicines and supplements should not be taken, unless, your physicians or nurse has advised you otherwise.

If you are unable to keep your appointment, please call the Medical Procedures Unit at (734) 936-9250 or toll-free at (877) 758-2626.

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#). Last Revised 01/2019