



Humalog® Junior KwikPen® (insulin lispro injection) 100 units/mL

Humalog Junior KwikPen Insulin Pen Parts:






Copyright © 2017, Eli Lilly and Company. All rights reserved.

Where to inject:

- Rotate injections between injection spots
 #1-4 in the diagram to the right:
 - 1. **Stomach**, except for the 2-inches around the belly button
 - 2. Top and outside of **thighs**, except in the bony area close to the knees
 - 3. Back of the **upper arms** (might work best if someone else is injecting)
 - 4. Top and outside of the **buttocks**
- Insulin is injected into the layer of fatty tissue under the skin, not into the muscle
- Do **not** inject into the same spot every time
- Do **not** inject into moles or scar



BruceBlaus [CC BY-SA 4.0], from Wikimedia Commons (Modified)

Step 1 Check The insulin



Prepare to inject:

- 1. If your Humalog Junior KwikPen pen is in the refrigerator, take it out at least 1-2 hours before use
- 2. Check the label to make sure you are using the right insulin pen
- 3. Check the expiration date
- 4. Remove the pen cap to check the insulin
 - Do **not** use insulin that is cloudy or not clear, or if it has any specks



Step 2 Attach The needle


- 5. Wipe the pen tip with an alcohol pad
- 6. Remove the seal on the needle cap
 - Do **not** share needles or insulin pens with anyone else
 - Do **not** re-use needles
- 7. Twist or push (based on the needle type) the needle straight onto the pen tip

Step 3 Perform an



For each injection:

- 1. Select a dose of 2 units
- 2. Take off the outer needle cap (save it) and inner needle cap (throw it away)



- 4. Press the button all the way in and make sure insulin comes out of the needle
 - Repeat up to two more times with the same needle if needed
 - If insulin does not come out after three times, change the needle and try again
 - If insulin still does not come out after changing the needle, the pen may be broken

Step 4 Select

The dose

Step 5

Inject The dose



- 5. Check that the dose counter shows "0" after the safety test
- 6. Turn the dose counter to the number of Humalog Junior KwikPen units that equals your dose
 - The whole unit numbers are printed on the dose counter, the half units are shown as lines between the numbers
 - Do **not** push the button while selecting your dose
- 7. Clean the injection site with an alcohol swab and wait for it to dry
- 8. Put the needle into the skin all the way
- 9. Press and hold the button to give the dose
- 10. Keep the button pressed and slowly count to 10 before taking the needle out of the skin
 - Do not rub the spot where you injected

Step 6 Remove And discard needle



- 11. Put the outer needle cap back on the needle and use it to remove the needle from the pen
 - Remove the needle from the pen after each dose
- 12. Place used needles in red sharps container or a hard plastic container (ex. an empty laundry detergent bottle)
 - Store the Humalog Junior KwikPen pen without a needle attached and with the pen cap on

Where to store Humalog Junior KwikPen:

- Unused pens should be kept in the refrigerator
- Open or in-use pens should be kept at room temperature for a maximum of 28 days
- After 28 days, Humalog Junior KwikPen pens should be thrown away in the trash even if there is leftover insulin
- Do not keep Humalog Junior KwikPen where there is direct heat or direct light
- If Humalog Junior KwikPen is frozen or overheated, throw it away

Top five tips for using your insulin:

- 1. Use Humalog Junior KwikPen prior to meals and snacks, as directed
- 2. Rotate your injections between the four injection spots listed above
- 3. Always use a new needle for each injection and clean the injection site before each dose
- 4. Slowly count to 10 after pressing the dose button
- 5. If insulin leaks after the injection, do **not** re-dose; be sure to keep the needle in the skin for at least 10 seconds next time

If you would like to watch a video on how to inject insulin using your Humalog Junior KwikPen, you can visit:

https://www.humalog.com/type-1-diabetes/#junior_kwikpen



Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by <u>Michigan Medicine</u> is licensed under a <u>Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License</u>. Last Revised 01/2019