


Levemir® FlexTouch® (insulin detemir injection) 100 units/mL


Levemir FlexTouch Insulin Pen Parts:



© 2018 Novo Nordisk All rights reserved. USA17INP03989 March 2018

Where to inject:

- Rotate injections between injection spots #1 4 in the diagram to the right:
 - 1. **Stomach**, except for the 2-inches around the belly button
 - 2. Top and outside of **thighs**, except in the bony area close to the knees
 - 3. Back of the **upper arms** (might work best if someone else is injecting)
 - 4. **Buttocks** is preferred in children for the best absorption possible
- Insulin is injected into the layer of fatty tissue under the skin, not into the muscle
- Do **not** inject into the same spot every time
- Do **not** inject into moles or scars



nsulin Injection Sites Back of the Body BruceBlaus [CC BY-SA 4.0], from Wikimedia Commons (Modified)

STEP 1 CHECK THE INSULIN



Prepare to inject:

- 1. If your Levemir FlexTouch pen is in the refrigerator, take it out at least 1-2 hours before use
- 2. Check the label to make sure you are using the right insulin pen
- 3. Check the expiration date
- 4. Remove the pen cap to check the insulin.
 - Do **not** use insulin that is cloudy or not clear, or if it has any specks

Department of Pharmacy/Pediatric Endocrinology

\bigcirc	5. Wipe the pen tip with an alcohol pad
	6. Remove the seal on the needle cap
ATTACH	• Do not share needles or insulin pens with anyone else
THE NEEDLE	• Do not re-use needles
	7. Twist or push (based on the needle type) the needle straight onto the
	pen tip
STEP 3	For each injection:
PERFORM	1. Select a dose of 2 units
AN AIR SHOT	2. Take off outer needle cap (save it)
	and inner needle cap (throw it away)
Part and Par	3. With the pen pointing up, tap the insulin to move the air bubbles to
	the top
4. Press the button all th	e way in and make sure insulin comes out of the needle
• Repeat up to two n	nore times with the same needle if needed
• If insulin does not	come out after three times, change the needle and try again
• If insulin still does	not come out after changing the needle, the pen may be broken
STEP 4	E Check that the does counter about "0" after the sefects test
SELECT	 Check that the dose counter shows "0" after the safety test Turn the dose counter to the number of Levemir FlexTouch units
THE DOSE	
Anterna B-	that equals your dose
	 Do not push the button while selecting your dose
	7 Chan the initial site with an alashal such and with family to der
STEP 5	7. Clean the injection site with an alcohol swab and wait for it to dry
STEP 5 INJECT	8. Put the needle into the skin all the way
STEP 5 INJECT THE DOSE	8. Put the needle into the skin all the way9. Press the button to give the dose
STEP 5 INJECT THE DOSE	8. Put the needle into the skin all the way9. Press the button to give the dose10.Keep the button pressed and slowly count to 10 before taking the
STEP 5 INJECT THE DOSE	8. Put the needle into the skin all the way9. Press the button to give the dose10.Keep the button pressed and slowly count to 10 before taking the needle out of the skin
STEP 5 INJECT THE DOSE	 8. Put the needle into the skin all the way 9. Press the button to give the dose 10. Keep the button pressed and slowly count to 10 before taking the needle out of the skin Do not rub the spot where you injected
STEP 6	 8. Put the needle into the skin all the way 9. Press the button to give the dose 10.Keep the button pressed and slowly count to 10 before taking the needle out of the skin Do not rub the spot where you injected 11.Put the outer needle cap back on the needle and use it to remove the
STEP 6 REMOVE	 8. Put the needle into the skin all the way 9. Press the button to give the dose 10.Keep the button pressed and slowly count to 10 before taking the needle out of the skin Do not rub the spot where you injected 11.Put the outer needle cap back on the needle and use it to remove the needle from the pen
STEP 6	 8. Put the needle into the skin all the way 9. Press the button to give the dose 10.Keep the button pressed and slowly count to 10 before taking the needle out of the skin Do not rub the spot where you injected 11.Put the outer needle cap back on the needle and use it to remove the needle from the pen Remove the needle from the pen after each dose
STEP 6 REMOVE AND DISCARD	 8. Put the needle into the skin all the way 9. Press the button to give the dose 10. Keep the button pressed and slowly count to 10 before taking the needle out of the skin Do not rub the spot where you injected 11. Put the outer needle cap back on the needle and use it to remove the needle from the pen Remove the needle from the pen after each dose 12. Used needles should be placed in red sharps container or a hard
STEP 6 REMOVE AND DISCARD	 8. Put the needle into the skin all the way 9. Press the button to give the dose 10. Keep the button pressed and slowly count to 10 before taking the needle out of the skin Do not rub the spot where you injected 11. Put the outer needle cap back on the needle and use it to remove the needle from the pen Remove the needle from the pen after each dose 12. Used needles should be placed in red sharps container or a hard plastic container (ex. empty laundry detergent bottle)
STEP 6 REMOVE AND DISCARD	 8. Put the needle into the skin all the way 9. Press the button to give the dose 10. Keep the button pressed and slowly count to 10 before taking the needle out of the skin Do not rub the spot where you injected 11. Put the outer needle cap back on the needle and use it to remove the needle from the pen Remove the needle from the pen after each dose 12. Used needles should be placed in red sharps container or a hard

Where to store Levemir FlexTouch:

- Unused pens should be kept in the refrigerator
- Open or in-use pens should be kept at room temperature for a maximum of 42 days
- After 42 days, Levemir FlexTouch pens should be thrown away in the trash even if there is leftover insulin
- Do **not** keep Levemir FlexTouch where there is direct heat or direct light
- If Levemir FlexTouch is frozen or overheated, throw it away

Top five tips for using your insulin:

- 1. Use Levemir FlexTouch once a day at the same time every day
- 2. Rotate your injections between the four injection spots listed above
- Always use a new needle for each injection and clean the injection site before each dose
- 4. Slowly count to 10 after pressing the dose button
- If insulin leaks after the injection, do not re-dose; be sure to keep the needle in the skin for at least 10 seconds next time

If you would like to watch a video on how to inject insulin using your Levemir FlexTouch, you can visit:

https://www.levemir.com/starting-on-levemir.html



Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by <u>Michigan Medicine</u> is licensed under a <u>Creative Commons Attribution</u>. <u>NonCommercial-ShareAlike 4.0 International Public License</u>. Last Revised 01/2019