



Uterine Fibroid Embolization Discharge Instructions

Symptoms to expect after your procedure typically include:

- Pelvic pain or cramping
- Nausea or vomiting
- Feeling tired
- Low-grade fever
- Poor appetite
- Flu-like symptoms
- Light spotting (can last up to 1 month)

These symptoms usually last 5-10 days and may be more severe in some women than others. Often, people feel better 1-2 days after the procedure and then experience additional episodes of pain and cramping. For this reason, we recommend taking your medications as prescribed for the first 3-4 days.

Special instructions

- It is a good idea to have a heating pad at home to apply to your abdomen.
- You may want to wear a light day maxi-pad if you have spotting.

When should I call for help?

Call Vascular and Interventional Radiology at (734) 936-4500, option 1, 5, if any of the following occur:

- Fever greater than 102.5 ° or fever associated with sweating and chills
- Abnormal foul-smelling vaginal discharge
- Heavy bright red vaginal bleeding, different from your normal menstrual flow
- Pain or burning with urination
- Pelvic pain unrelieved by pain medication, or pain which lasts longer than two weeks

What are my activity instructions?

- You may remove the band-aid from your groin after 24 hours and shower
- Avoid strenuous physical activity for 48 hours
- Activity is not restricted after 48 hours. After the first 2-3 days, the way you feel is the best indicator of what you should and should not do
- There are no restrictions on resuming sexual activities
- If you are taking pain medication, do not drive or operate machinery

What are my diet instructions?

- You may resume your previous diet as tolerated.
- You may want to add more fiber to your diet in the form of fruits and vegetables to prevent constipation while taking pain medications.

What are my medication instructions?

- Follow the medication regimen closely for the first 3-4 days.
- You will receive prescriptions at the time of discharge for pain medication. Usually Motrin, Colace (a stool softener), and Zofran (for nausea).

What follow-up care will I receive?

Follow-up appointments are scheduled at two weeks and six months.

If for some reason you do not receive an appointment, please call our office at (734) 936-4500, option 1, 5, to schedule your two-week follow-up appointment.

Disclaimer: This document contains information and/or instructional materials developed by Michigan Medicine for the typical patient with your condition. It may include links to online content that was not created by Michigan Medicine and for which Michigan Medicine does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Patient Education by [Michigan Medicine](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#). Last Revised 1/2023